

Personnelle Cold and Flu-in-One Extra Strength

Outer product packaging:



Warnings on the inside of the outer product packaging:

CAUTION: Consult a doctor if sore throat pain persists for more than 2 days, if fever persists for more than 3 days or symptoms more than 5 days, if cough worsens, tends to recur, or is accompanied by high fever ($>38^{\circ}\text{C}$) or the production of yellow/green phlegm, or if you develop nervousness, dizziness, sleeplessness or allergic reactions such as wheezing, rash or itching. Consult a doctor or pharmacist prior to use if you use any other medications, including natural health products, prescription drugs, salicylates or other pain and fever relief medications (nonsteroidal anti-inflammatory drugs (NSAIDs)), if you have chronic alcoholism, serious kidney or liver disease, heart or thyroid disease, chronic lung disease, high blood pressure, diabetes, glaucoma, difficulty in urination due to an enlargement of the prostate gland or if you are pregnant or nursing. Not recommended for patients with asthma, shortness of breath or persistent chronic cough unless directed by a physician. Avoid alcohol.

IN CASE OF OVERDOSE: Call a doctor or Poison Control Centre immediately, even if you do not notice any signs or symptoms. Within the first 24 hours, you may experience increased sweating, nausea, vomiting, stomach pain and loss of appetite.

Personnelle Flu Relief

Outer product packaging:



Warnings on the inside of the outer product packaging:

ADDITIONAL INFORMATION FOR THE CONSUMER

WARNING: Talk to a doctor if fever persists for more than 3 days or symptoms last more than 5 days, if cough worsens, tends to recur or is accompanied by high fever ($>38^{\circ}\text{C}$) or the production of yellow/green phlegm, or if you develop nervousness, dizziness, sleeplessness, or allergic reactions such as wheezing, rash or itching. Consult a doctor or pharmacist prior to use if you use any other medications, including natural health products, prescription drugs, salicylates or other pain and fever relief medications (nonsteroidal anti-inflammatory drugs (NSAIDs)), if you have chronic alcoholism, serious kidney or liver disease, heart or thyroid disease, asthma, shortness of breath, persistent chronic cough, chronic lung disease, high blood pressure, diabetes, glaucoma, or difficulty in urination due to an enlargement of the prostate gland or if you are pregnant or nursing. Nighttime caplets may cause drowsiness or excitability. Avoid alcohol, driving or other activities requiring alertness. Not to be used by elderly patients who experience confusion at nighttime. These drugs may produce excitation rather than sedation in the elderly. Therefore, they should be avoided in this age group.

IN CASE OF OVERDOSE: Call a doctor or Poison Control Centre immediately, even if you do not notice any signs or symptoms. Within the first 24 hours, you may experience increased sweating, nausea, vomiting, stomach pain and loss of appetite.